

# Expanded Inverse-Sandwich Complexes of Lanthanum Borides: $\text{La}_2\text{B}_{10}^-$ and $\text{La}_2\text{B}_{11}^-$

Published as part of *The Journal of Physical Chemistry virtual special issue "Alexander Boldyrev Festschrift"*.

Zhi-Yu Jiang,<sup>#</sup> Teng-Teng Chen,<sup>#</sup> Wei-Jia Chen, Wan-Lu Li,<sup>\*</sup> Jun Li,<sup>\*</sup> and Lai-Sheng Wang<sup>\*</sup>



Cite This: *J. Phys. Chem. A* 2021, 125, 2622–2630



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

**ABSTRACT:** Inverse-sandwich structures have been observed recently for dilanthanide boride clusters, in which two Ln atoms sandwich a monocyclic  $\text{B}_x$  ring for  $x = 7-9$ . An interesting question is if larger  $\text{B}_x$  rings are possible to form such inverse-sandwich clusters. Here we address this question by investigating  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$  using photoelectron spectroscopy and *ab initio* quantum chemical calculations. Photoelectron spectra of  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$  show complicated, but well-resolved, spectral features that are used to compare with theoretical calculations. We have found that global minimum structures of the two clusters are based on the octa-boron ring. The global minimum of  $\text{La}_2\text{B}_{10}^-$  consists of two chiral enantiomers with  $C_1$  symmetry, which can be viewed as adding a  $\text{B}_2$  unit off-plane to the  $\text{B}_8$  ring, whereas that of  $\text{La}_2\text{B}_{11}^-$  can be viewed as adding a  $\text{B}_3$  unit in-plane to the  $\text{B}_8$  ring in a second coordination shell. Chemical bonding analyses reveal localized B–B bonds on the edge of the clusters and delocalized bonds in the expanded boron frameworks. The interactions between the La atoms and the boron frameworks include the unique  $(d-p)\delta$  bonding, which was found to be the key for inverse-sandwich complexes with monocyclic boron rings. The current study confirms that the largest monocyclic boron ring to form the inverse-sandwich structures is  $\text{B}_9$  and provide insights into the structural evolutions of larger lanthanide boride clusters.



## 1. INTRODUCTION

Boron forms a wide range of important boride materials, from the superconducting  $\text{MgB}_2$  and superhard transition-metal borides to borides with ultrahigh thermal conductivity.<sup>1–6</sup> Lanthanide (Ln) borides constitute another interesting class of boride materials with unique magnetic, optical, superconducting, mechanical, and thermoelectric properties.<sup>7–10</sup> Over the past two decades, the structures and chemical bonding of size-selected bare boron clusters have been studied by joint experimental and theoretical methods,<sup>11–15</sup> discovering planar clusters, nanotubular structures, graphene-like borophenes, and fullerene-like borospherenes.<sup>15–22</sup> Transition-metal-doped boron clusters have led to a new direction of boron nanoclusters, giving rise to new structures and bonding, such as the metal-centered aromatic borometallic wheels<sup>23–25</sup> and tubular metal-centered drums.<sup>26–29</sup> However, lanthanide boride clusters have been rarely explored until recently,<sup>30–40</sup> despite the importance of bulk lanthanide boride materials.

The  $\text{PrB}_7^-$  cluster represents the first lanthanide boride cluster characterized by photoelectron spectroscopy (PES) and quantum chemistry calculations.<sup>32</sup> It was found to have a half-sandwich structure with the Pr atom coordinated to a doubly aromatic bowl-shaped  $\eta\text{-B}_7^{3-}$  unit. A series of small lanthanide boride clusters,  $\text{SmB}_6^-$ ,  $\text{CeB}_6^-$ ,  $\text{PrB}_3^-$ , and  $\text{PrB}_4^-$ ,<sup>33–35</sup> have been reported, which all have planar structures with the Ln atom on the edge. Most recently, larger lanthanide boride clusters have been studied, uncovering an inverse triple-decker


and an interesting cage-like spherical trihedral metallo-borospherene.<sup>36,37</sup> Among all the lanthanide boride clusters, photoelectron spectra of the dilanthanide octaboron clusters ( $\text{Ln}_2\text{B}_8^-$ , Ln = La, Pr, Tb) were observed to be relatively simple and these clusters were found to consist of unprecedented inverse-sandwich structures with  $D_{8h}$  symmetry,<sup>38</sup> in which the two Ln atoms sandwiching the doubly aromatic  $\text{B}_8$  monocyclic ring with strong  $(d-p)\pi$  and  $(d-p)\delta$  bonding between the Ln atoms and the  $\text{B}_8$  ring. Subsequently, two more dilanthanide inverse-sandwich complexes,  $\text{La}_2\text{B}_7^-$  and  $\text{La}_2\text{B}_9^-$ , were characterized, suggesting that there exists a class of such inverse-sandwich structures with different boron ring sizes.<sup>39</sup> An interesting question is can inverse-sandwiches be formed with larger  $\text{B}_x$  rings? What are the structures of the larger dilanthanide boride clusters ( $\text{La}_2\text{B}_x^-$ ,  $x \geq 10$ )?

In this article we aim to answer these questions and report an investigation of  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$  using PES and quantum calculations. Our global-minimum structural searches show that the most stable structure of  $\text{La}_2\text{B}_{10}^-$  consists of a pair

Received: February 7, 2021

Revised: March 10, 2021

Published: March 19, 2021



of chiral enantiomers ( $C_1$ ,  $^1A$ ), which can be viewed as adding a  $B_2$  unit off-plane to a  $La_2B_8$  inverse sandwich. The global minimum of  $La_2B_{11}^-$  is also based on the  $La_2B_8$  inverse-sandwich structure, with the three extra B atoms being added to the  $B_8$  ring in-plane by forming a second coordination shell and a perfectly planar expanded  $B_{11}$  framework. Chemical bonding analyses show localized B–B bonds on the edge and multicentered B–B bonds on the expanded boron frameworks. The expanded inverse sandwiches also feature the (d–p) $\delta$  bonding between the La atoms and the boron frameworks. The current study enriches the family of lanthanide inverse-sandwich clusters, providing insight into the formation and stability of larger lanthanide boride clusters.

## 2. METHODS

**2.1. Photoelectron Spectroscopy.** The experiments were carried out using a magnetic-bottle PES apparatus, coupled with a time-of-flight mass spectrometer and a laser vaporization cluster source, details of which have been published before.<sup>12</sup> Briefly, the lanthanide boride clusters were produced by laser vaporization of a disc target made from a mixed powder of La (Alfa Aesar, 99.9% purity) and B (Alfa Aesar, 96%  $^{11}B$  enriched, 99.9% elemental purity) with a 5/2 La/B mass ratio. The laser-induced plasma was quenched by a He carrier gas seeded with 5% Ar inside the nozzle to initiate nucleation. The nascent clusters were entrained by the carrier gas and underwent a supersonic expansion to form a cold cluster beam. After passing a skimmer, negatively charged clusters were extracted from the collimated cluster beam and analyzed by a time-of-flight mass spectrometer. A range of  $La_2B_x^-$  clusters were formed in the source. The  $La_2B_{10}^-$  and  $La_2B_{11}^-$  clusters of current interest were mass-selected and decelerated before being photodetached by the 193 nm radiation from an ArF excimer laser. As discussed previously,<sup>12</sup> the cluster temperatures depended on the resident time inside the nozzle. We usually selected clusters that had the longest resident time because they were colder. Photoelectrons were collected at nearly 100% efficiency by the magnetic bottle and analyzed in a 3.5 m long electron flight tube. Photoelectron spectra were calibrated using the known spectra of the  $Au^-$  and  $Bi^-$  atomic anions. The electron kinetic energy (KE) resolution of the PES apparatus ( $\Delta KE/KE$ ) was around 2.5%, that is,  $\sim 25$  meV for 1 eV electrons.


**2.2. Computational Methods.** Unbiased structural searches for the global minima of  $La_2B_{10}^-$  and  $La_2B_{11}^-$  with different spin multiplicities were carried out using the TGMIn 2.0 package,<sup>41–43</sup> combined with energy calculations using the ADF 2017.114 software.<sup>44–46</sup> In the structure search, the Perdew–Burke–Ernzerhof (PBE) generalized gradient approximation (GGA) density functional<sup>47</sup> and Slater-type basis sets with the triple- $\zeta$  plus one polarization function (TZP) were used.<sup>48</sup> The scalar relativistic effects were considered by the zero-order-regular approximation (ZORA).<sup>49</sup> All local minima were verified by calculations of the harmonic vibrational frequencies. Low-lying isomers were subsequently reoptimized using the hybrid PBE0 functional in Gaussian 16.<sup>50</sup> In the reoptimization calculations, the cc-pVTZ valence triple- $\zeta$  basis set was used for boron<sup>51</sup> and the Stuttgart energy-consistent relativistic pseudopotential ECP28MWB(La) with the corresponding ECP28MWB\_ANO basis set for La.<sup>52,53</sup>

Photodetachment transitions were computed approximately using the  $\Delta SCF$ -TDDFT approach.<sup>54</sup> The first vertical detachment energy (VDE<sub>1</sub>) was calculated as the energy

difference between the anion and the neutral at the anion geometry. Vertical excitation energies of the neutral species calculated using the time-dependent DFT method and the SAOP model (TD–SAOP) were added to VDE<sub>1</sub> to yield the higher VDEs.<sup>55</sup> The obtained VDEs were fitted with unit-area Gaussians of 0.15 eV width to produce a simulated spectrum. Chemical bonding was analyzed using the adaptive natural density partitioning (AdNDP)<sup>56</sup> using the PBE0/cc-pVTZ/ECP28MWB method based on the geometry at the same level from Gaussian 16. The charges of the atoms were calculated using various partition schemes.<sup>57–61</sup>

## 3. RESULTS


**3.1. Photoelectron Spectroscopy.** The photoelectron spectra of  $La_2B_{10}^-$  and  $La_2B_{11}^-$  at 193 nm (6.424 eV) are shown in Figures 1 and 2, respectively. The PES bands are



**Figure 1.** (a) Photoelectron spectrum of  $La_2B_{10}^-$  at 193 nm (6.424 eV) and (b) simulated spectrum of the global minimum isomer I/II at the TD-SAOP/TZP level.

labeled with letters (X, A, B, ...), and the VDEs measured from the maxima of all the observed bands are given in Tables 1 and 2 for  $La_2B_{10}^-$  and  $La_2B_{11}^-$ , respectively, where they are compared with the theoretical results.

The 193 nm spectrum of  $La_2B_{10}^-$  shows seven well-resolved bands, labeled as X, A–F in Figure 1a. The lowest-binding-energy band (X) gives rise to the first VDE of 1.69 eV. The first adiabatic detachment energy (ADE) is estimated from the onset of band X as 1.48 eV, which also represents the electron affinity (EA) of the corresponding neutral  $La_2B_{10}$ . The next band A is observed at 2.14 eV followed by band B at 2.60 eV. Two intense bands, C and D, are observed at 3.23 and 3.77 eV, respectively. A broad and weaker band E appears at 4.15 eV, followed by a small energy gap and a broad band F at 5.01 eV. Beyond 5.5 eV, the signal-to-noise ratio becomes poor and a band G at  $\sim 6$  eV is tentatively labeled for the sake of discussion. Unlike the previously reported spectra of the inverse-sandwich complexes  $La_2B_x^-$  ( $x = 7–9$ ),<sup>38,39</sup> the spectrum of  $La_2B_{10}^-$  displays complicated and broad features, indicating a low symmetry structure.



**Figure 2.** (a) Photoelectron spectra of  $\text{La}_2\text{B}_{11}^-$  at 193 nm (6.424 eV) and (b) the simulated spectrum of the global minimum structure at the TD-SAOP/TZP level.

The 193 nm spectrum of  $\text{La}_2\text{B}_{11}^-$  appears simpler than that of  $\text{La}_2\text{B}_{10}^-$ , displaying five major bands, as shown in Figure 2a. The most intense band X yields the first VDE at 2.61 eV, but its high relative intensity and large spectral width suggests that it likely contains multiple detachment channels. The first ADE, i.e., the EA of  $\text{La}_2\text{B}_{11}$ , is estimated from the onset of band X to be 2.27 eV, which is much higher than that of  $\text{La}_2\text{B}_{10}$ . Following a large energy gap, a well-resolved band A is observed at 3.63 eV. The next spectral feature is quite broad

and three overlapping bands are discernible: band B at 4.06 eV, band C at 4.21 eV, and band D at 4.49 eV. A relatively weak band E at 4.98 eV is well resolved, following a small energy gap. A broad and strong band F at  $\sim 6.0$  eV is observed at the high binding energy side, which probably contains multiple detachment channels.

**3.2. Global Minima of  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$ .** The optimized structures within 50 kcal/mol of the global minima of  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$  are presented in Figures S1 and S2, respectively. All the structures were determined to be true minima without imaginary frequencies. We found that the global minimum of  $\text{La}_2\text{B}_{10}^-$  consists of two degenerate expanded inverse-sandwich structures, I and II ( $C_1$ ,  $^2A$ ), as shown in Figure 3. Interestingly, these two structures are a pair of chiral enantiomers. The top and bottom views for each enantiomer are shown in Figure 3, and the detailed bond lengths of each structure are given in Figure S3. These structures can be viewed as adding a  $\text{B}_2$  unit *parallelly* above the  $\text{B}_8$ -ring plane of a  $\text{La}_2\text{B}_8^-$  inverse sandwich. The  $\text{B}_2$  unit distorts the local  $\text{B}_8$  ring structure, creating a low-symmetry cluster ( $C_1$ ) in which one La is coordinated by a  $\text{B}_8$  ring on one side and the other La is coordinated by a  $\text{B}_9$  ring on the other side of the inverse sandwich. The next low-lying isomer III ( $C_{2v}$ ,  $^2A_1$ ; Figure S4) is 3.24 kcal/mol at the PBE level (3.75 kcal/mol at the PBE0 level) higher in energy above the global minimum (Figure S1). Isomer III can be viewed as inserting a  $\text{B}_2$  unit *perpendicularly* into the  $\text{B}_8$ -ring of a  $\text{La}_2\text{B}_8^-$  inverse sandwich. This isomer has a higher symmetry ( $C_{2v}$ ) and each La atom is coordinated by a  $\text{B}_9$  ring.

The global minimum of  $\text{La}_2\text{B}_{11}^-$  was found to have a closed-shell  $C_{2v}$  ( $^1A_1$ ) inverse-sandwich structure, as shown in Figure 4. The structure can be viewed as a  $\text{La}_2\text{B}_8^-$  inverse sandwich, where the additional three B atoms are added to the same plane as the  $\text{B}_8$  ring in a second coordination shell. The

**Table 1. Experimental Vertical Detachment Energies (VDEs) of  $\text{La}_2\text{B}_{10}^-$  in Comparison with Theoretical VDEs Computed for the Global Minimum Structure I/II ( $C_1$ ,  $^2A$ ) at the TD-SAOP/TZP Level (All Energies in eV)**


feature	VDE (exp) <sup>a</sup>	final state and electronic configuration xs(SAOP)	VDE (theo)
X	1.69	$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^2 82a^2 83a^0 \}$	1.64
A	2.14	$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^2 82a^1 83a^1 \}$	2.06
		$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^2 82a^1 83a^1 \}$	2.24
B	2.60	$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	2.50
		$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	2.64
C	3.23	$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^1 81a^1 82a^2 83a^1 \}$	3.12
		$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^1 80a^1 81a^1 82a^2 83a^1 \}$	3.16
		$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^1 81a^1 82a^2 83a^1 \}$	3.35
D	3.77	$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^1 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	3.44
		$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^1 79a^1 80a^2 81a^1 82a^2 83a^1 \}$	3.48
		$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^1 78a^1 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	3.60
		$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^2 78a^1 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	3.61
E	4.15	$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^2 77a^1 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	3.88
		$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^1 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	4.05
		$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^2 76a^1 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	4.24
F	5.01	$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^1 76a^2 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	4.61
		$^3A \{ \dots 0.71a^2 72a^2 73a^2 74a^1 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	4.64
		$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^2 75a^1 76a^2 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	4.75
		$^1A \{ \dots 0.71a^2 72a^2 73a^2 74a^1 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	4.85
		$^3A \{ \dots 0.71a^2 72a^2 73a^1 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	4.94
G	$\sim 6$	$^3A \{ \dots 0.71a^2 72a^1 73a^2 74a^2 75a^2 76a^2 77a^2 78a^2 79a^2 80a^2 81a^1 82a^2 83a^1 \}$	5.94

<sup>a</sup>The experimental uncertainty was estimated to be  $\pm 0.03$  eV.


**Table 2.** Experimental Vertical Detachment Energies (VDEs) of  $\text{La}_2\text{B}_{11}^-$  in Comparison with Theoretical VDEs Computed from the Global Minimum Structure ( $C_{2v}$ ,  $^1A_1$ ) at the TD-SAOP/TZP Level (All Energies in eV)

feature	VDE (exp) <sup>a</sup>	final state and electronic configuration (SAOP)	VDE (theo)
X	2.61	$^2B_1 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	2.32
		$^2B_2 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	2.42
		$^2A_2 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	2.47
A	3.63	$^2A_2 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	3.37
B	4.06	$^2A_1 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	3.62
C	4.21	$^2B_2 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	3.77
D	4.49	$^2A_1 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	3.97
E	4.98	$^2B_1 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	4.32
F	~6.0	$^2B_1 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	4.58
		$^2B_2 \{ \dots 0.32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	5.45
		$^2A_1 \{ \dots 32a_1^2 20b_2^2 16b_1^2 17b_1^2 18b_1^2 33a_1^2 21b_2^2 34a_1^2 9a_2^2 10a_2^2 22b_2^2 19b_1^2 \}$	5.59

<sup>a</sup>The experimental uncertainty was estimated to be  $\pm 0.03$  eV.



**Figure 3.** Top and bottom views of the two enantiomers of the chiral global minimum of  $\text{La}_2\text{B}_{10}^-$ .



**Figure 4.** Top and side views of the optimized global minimum structure of  $\text{La}_2\text{B}_{11}^-$ .

detailed bond lengths of the global minimum are shown in Figure S5. The nearest low-lying isomer II (Figure S2) has  $C_1$  symmetry and it is 8.86 kcal/mol (at the PBE/TZP level) and 11.96 kcal/mol (at the PBE0/TZP level) higher in energy than the global minimum, suggesting the high stability of the  $C_{2v}$ ,  $\text{La}_2\text{B}_{11}^-$  inverse-sandwich structure.

To validate the global minimum structures of  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$ , we have calculated their VDEs, as compared with the experimental data in Tables 1 and 2, as well as in Figures 1b and 2b, respectively, at the TD-SAOP/TZP level. Because isomer III of  $\text{La}_2\text{B}_{10}^-$  is relatively low-lying, we also computed its VDEs (Table S1) and compared its simulated spectrum with the experiment (Figure S6). The VDE<sub>1</sub> and ADE values for the chiral global minimum of  $\text{La}_2\text{B}_{10}^-$  and the  $C_{2v}$  global minimum of  $\text{La}_2\text{B}_{11}^-$  were computed at different levels of theory, as summarized in Table S2. The atomic charges for isomers I/II and III of  $\text{La}_2\text{B}_{10}^-$  and that of isomer I of  $\text{La}_2\text{B}_{11}^-$  are given in Tables S3 and S4, respectively. The coordinates for isomers I/II and III of  $\text{La}_2\text{B}_{10}^-$  and that of isomer I of  $\text{La}_2\text{B}_{11}^-$  at the PBE0 level are given in Table S5. Except for their structures, the chiral global minimum isomers I/II of  $\text{La}_2\text{B}_{10}^-$  have identical electronic structure and chemical bonding properties.

## 4. DISCUSSION

**4.1. Comparison between Experiment and Theory.** As shown in Table S2, both PBE and PBE0 functionals in general gave similar VDE<sub>1</sub> and ADE values for the global minima of  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$ , both in good agreement with the experimental data. Each computed VDE at the TD-SAOP/TZP level for the global minima is fitted with a Gaussian function of unit area and a width of 0.15 eV to produce the simulated spectra, which are compared with the experimental data in Figures 1b and 2b for  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$ , respectively.

**4.1.1.  $\text{La}_2\text{B}_{10}^-$ .** The global minimum chiral structure of  $\text{La}_2\text{B}_{10}^-$  has a doublet ground state ( $^2A$ ) with a single electron in the 83a orbital (SOMO, Figure S7a). The computed VDE<sub>1</sub>/ADE of 1.67/1.48 eV at the PBE/TZP level (Table S2) by removing the electron from the SOMO are in excellent agreement with the experimental data of 1.69/1.48 eV. The removal of an electron from the HOMO (82a, Figure S7a) gives rise to two final spin states, one triplet state with a computed VDE of 2.06 eV and one singlet state with a computed VDE of 2.24 eV; both are consistent with the observed band A at 2.14 eV (Table 1). Similarly, detachment from the HOMO-1 (81a) yields a triplet state with a computed VDE at 2.50 eV and a singlet state at 2.64 eV, both matching well with the observed band B at 2.60 eV (Table 1). Band C at 3.23 eV and band D at 3.77 eV contain multiple




Figure 5. AdNDP analyses of the global minimum  $C_1$  chiral structure of  $La_2B_{10}^-$  at the PBE0/VTZ level.





Figure 6. AdNDP analyses of the global minimum  $C_{2v}$  structure of  $La_2B_{11}^-$  at the PBE0/VTZ level.

detachment channels from the HOMO-2 (80a), HOMO-3 (79a), HOMO-4 (78a), and HOMO-5 (77a), while band E at 4.15 eV agrees well with the computed VDEs from the HOMO-6 (76a). Band F is quite broad and in good agreement with detachment channels from the HOMO-7 (75a), HOMO-8 (74a) and HOMO-9 (73a), while the tentative band G can be explained by detachment from the 72a orbital (Table 1). The simulated spectrum of the  $C_1$  chiral structure of  $La_2B_{10}^-$  reproduces well the experimental features (Figure 1b). On the other hand, the low-lying isomer III gives rises to a relatively simple simulated spectral pattern due to its higher symmetry, which does not agree with the experiment, as shown in Figure S6. Thus, the good agreement between the

experimental and theoretical data provides strong evidence for the chiral  $C_1$  structure as the global minimum of  $La_2B_{10}^-$ .

**4.1.2.  $La_2B_{11}^-$ .** The global minimum of  $La_2B_{11}^-$  (Figure 4) is closed shell ( $^1A_1$ ). Thus, electron detachment from each occupied orbital yields a single doublet final state (Table 2). The computed VDE<sub>1</sub>/ADE values of 2.34/2.23 eV at the PBE/TZP level (Table S2) are in reasonable agreement with the experimental values of 2.67/2.27 eV. The larger discrepancy between the computed VDE<sub>1</sub> and the experimental VDE<sub>1</sub> is due to the fact that the first real detachment channel at a lower VDE is not resolved in the broad band X in the 193 nm spectrum, as can be seen by the shoulder on the lower binding energy side of band X (Figure 2a). The X band is consistent with the first three detachment channels from the HOMO



**Figure 7.** MO bonding scheme of the  $C_{2v}$  global minimum of  $La_2B_{11}^-$  at the level of PBE0/TZP, illustrating the bonding interactions between the  $La \cdots La$  and  $B_{11}^-$  framework (isovalue = 0.03 au).

( $19b_1$ ), HOMO-1 ( $22b_2$ ), and HOMO-2 ( $10a_2$ ) (Table 2), which give rise to closely spaced VDEs. Electron removal from the HOMO-3 ( $9a_2$ ), HOMO-4 ( $34a_1$ ), HOMO-5 ( $21b_2$ ), and HOMO-6 ( $33a_1$ ) results in four closely spaced detachment channels with computed VDEs ranging from 3.37 to 3.97 eV, which agree well with bands A, B, C, D, respectively (Table 2). These are followed by two detachment channels from the HOMO-7 ( $18b_1$ ) and HOMO-8 ( $17b_1$ ) with computed VDEs of 4.32 and 4.58 eV, which can be assigned to the broad band E. The next three detachment channels from the HOMO-9 ( $16b_1$ ), HOMO-10 ( $20b_2$ ), and HOMO-11 ( $32a_1$ ) give rise to three closely spaced VDEs from 5.45 to 5.59 eV (Table 2), consistent with the broad band F at the high binding energy side (Figure 2). The overall spectral pattern of the simulated spectrum agrees well with the experiment (Figure 2), confirming the inverse-sandwich  $C_{2v}$  structure to be the global minimum of  $La_2B_{11}^-$ .

**4.2. Chemical Bonding Analyses.** The low-symmetry structures, in particular the  $C_1$  global minimum of  $La_2B_{10}^-$ , make it a challenge to decipher the chemical bonding in the two large dilanthanum boride clusters. Because of the electron deficiency of the B atom, multicenter bonding is prevalent in boron clusters.<sup>62–64</sup> In general, for the dilanthanide inverse sandwiches the B 2s and in-plane 2p orbitals form the  $\sigma$ -bonding network, while the two Ln atoms form symmetric and antisymmetric  $\sigma$  ( $d_z^2$ ),  $\pi$  ( $d_{xz}$ ,  $d_{yz}$ ) and  $\delta$  ( $d_{x^2-y^2}$ ,  $d_{xy}$ ) type d-orbital pairs that can interact with the out-of-plane B 2p orbitals of the ring via  $(d-p)\sigma$ ,  $(d-p)\pi$ , and  $(d-p)\delta$  bonds.<sup>38–40</sup> The canonical Kohn–Sham molecular orbitals of isomers I/II and III of  $La_2B_{10}^-$  and isomer I of  $La_2B_{11}^-$  are

shown in Figure S7. To better understand the chemical bonding and stability of the global minima of  $La_2B_{10}^-$  and  $La_2B_{11}^-$ , we performed AdNDP analyses,<sup>56</sup> as shown in Figures 5 and 6, respectively. Because of the complexity of the bonding, we also did AdNDP analyses for the  $B_{10}^-$  and  $B_{11}^-$  frameworks or ligands without the La atoms, as shown in Figures S8 and S9.

**4.2.1.  $La_2B_{10}^-$ .** The AdNDP result for the  $B_{10}^-$  framework (Figure S8a) of the chiral global minimum of  $La_2B_{10}^-$  exhibit nine B–B two-center two-electron (2c–2e) bonds on the periphery, three 5c–2e delocalized bonds over the expanded part of the  $B_{10}$  framework. The remaining 7 electrons form four totally delocalized  $\sigma$  and  $\pi$  bonds over the entire  $B_{10}$  framework, which have orbital interactions with the two La atoms. Indeed, upon coordination with the  $La \cdots La$  atoms in the chiral  $La_2B_{10}^-$ , the nine 2c–2e bonds and the three 5c–2e bonds are largely unchanged (Figure 5). Six 12c–2e bonds and one 12c–1e bond are observed, suggesting that three unoccupied MOs in the  $B_{10}$  framework are stabilized significantly via metal–ligand interactions, so that the  $B_{10}^-$  ligand can be viewed to have gained 6 electrons in  $La_2B_{10}^-$ . Since the extra unpaired electron remains largely in a La-based orbital, the two La atoms can be viewed as having oxidation states of La(III) and La(II) in  $La_2B_{10}^-$ . The charge transfer from the  $La \cdots La$  atoms to the  $B_{10}^-$  framework is consistent with the charge population analyses (Table S3). The six 12c–2e bonds are formed from the interactions between the  $La \cdots La$   $5d_\sigma$ ,  $5d_\pi$ , and  $5d_\delta$  orbitals and the  $B_{10}^-$  ligand. The two 12c–2e ( $d-p$ ) $\delta$  bonds due to the interactions of the La  $5d_\delta$  orbitals and the  $\pi$  orbitals of the boron framework are particularly

interesting. Similar (d-p) $\delta$  bonds have been shown previously to play critical roles in stabilizing the monocyclic inverse-sandwich structures in  $\text{La}_2\text{B}_x^-$  ( $x = 7-9$ ).<sup>38-40</sup>

The AdNDP results for the low-lying  $C_{2v}$  isomer III of  $\text{La}_2\text{B}_{10}^-$  (Figure S10 and Figure S8b for the  $\text{B}_{10}^-$  ligand) are similar to that of the chiral global minimum, but the bonding is easier to visualize due to the higher symmetry of the  $C_{2v}$  isomer. One notable difference is the number of 2c-2e B-B bonds on the periphery of the two structures: nine in the  $C_1$  global minimum vs seven in the  $C_{2v}$  isomer. Presumably the more B-B 2c-2e  $\sigma$ -bonds in the  $C_1$  structures makes it more stable because the localized B-B bonds are stronger bonds than the multicentered bonds.

**4.2.2.  $\text{La}_2\text{B}_{11}^-$ .** The higher symmetry of the  $\text{La}_2\text{B}_{11}^-$  global minimum and its planar boron framework make it easier to visualize its bonding. The AdNDP analyses of the planar  $\text{B}_{11}^-$  framework (Figure S9) reveal ten 2c-2e B-B  $\sigma$  bonds on the periphery, three 6c-2e bonds (one  $\pi$  bond and two  $\sigma$  bonds) involving the second and first coordination shells, and four totally delocalized 11c-2e bonds (two  $\pi$  bonds and two  $\sigma$  bonds). The three  $\pi$  bonds fulfill the  $4N + 2$  Hückel rule, indicating that the  $\text{B}_{11}^-$  ligand is  $\pi$ -aromatic. Upon complexation to the two La...La atoms, the 2c-2e bonds and the 6c-2e bonds essentially remain the same while the 11c-2e bonds interact with the La atoms, as shown in Figure 6. There are seven delocalized 13c-2e bonds representing interactions between the La...La  $5d_{\sigma}$ ,  $5d_{\pi}$ , and  $5d_{\delta}$  orbitals and the  $\text{B}_{11}^-$  ligand. The seven 13c-2e bonds suggest that three unoccupied MOs of the  $\text{B}_{11}^-$  ligands are also involved in bonding with the La...La atoms and are significantly stabilized, as shown in Figure 7. Two of these MOs are  $\pi$  MOs ( $3b_2$  and  $2a_2$  in Figure 7), which form the two 13c-2e (d-p) $\delta$  bonds; one of these MOs ( $11b_1$  in Figure 7) forms a 13c-2e (d-p) $\pi$  bond. Thus, the  $\text{B}_{11}^-$  ligand can be viewed as gaining six electrons upon complexation with the two La...La atoms, indicating that the La atoms are in their stable La(III) oxidation state in  $\text{La}_2\text{B}_{11}^-$ , consistent with charge population analyses (Table S4). Figure 7 also shows that the  $\text{La}_2\text{B}_{11}^-$  cluster has an extremely stable closed-shell electron configuration with 40 valence electrons and a large HOMO-LUMO gap. Such a stable electronic structure is evidenced by the very high ADE of  $\text{La}_2\text{B}_{11}^-$  (2.27 eV) relative to that of  $\text{La}_2\text{B}_{10}^-$  (1.48 eV). In fact, the bonding in  $\text{La}_2\text{B}_{11}^-$  closely resemble that in the monocyclic  $\text{La}_2\text{B}_9^-$  system,<sup>39,40</sup> again confirming the importance of the (d-p) $\delta$  bonds in stabilizing the inverse-sandwich structure.

**4.3. Comparison with the Monocyclic  $\text{La}_2\text{B}_x^-$  ( $x = 7-9$ ) Systems.** The structures of  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$  confirm that for dilanthanide boride clusters ( $\text{Ln}_2\text{B}_x^-$ ) the largest monocyclic ring ( $\text{B}_x$ ) to form the inverse-sandwich structure in the 9-membered ring in  $\text{Ln}_2\text{B}_9^-$ .<sup>39,40</sup> Clearly, larger rings would reduce the orbital overlaps between the Ln...Ln atoms and the boron rings. Thus, for  $\text{Ln}_2\text{B}_x^-$  clusters with  $x > 9$ , the additional boron atoms do not expand the ring size. Although  $\text{La}_2\text{B}_9^-$  has a higher stability than  $\text{La}_2\text{B}_8^-$  due to the fully occupied (d-p) $\delta$  bonding orbitals,<sup>39,40</sup> the  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$  clusters are mainly based on a  $\text{B}_8$ -ring system. This can be understood readily from the fact that the  $\text{B}_8$  ring size in the  $\text{B}_{11}$  framework in  $\text{La}_2\text{B}_{11}^-$  is larger than the bare monocyclic  $\text{B}_8$  ring in the  $\text{La}_2\text{B}_8^-$  inverse sandwich because the B-B bond lengths for the two B atoms that are bonded to the second coordination shell are longer (Figure S5). This is consistent with the fact that the  $\text{B}_9$  monocyclic ring is the largest size to form inverse sandwiches in dilanthanide boride

clusters. It is conceivable that larger ring sizes may be possible for diactinide boride inverse-sandwich clusters due to the larger sizes of the actinide elements.

## 5. CONCLUSIONS

In conclusion, we report a joint photoelectron spectroscopy and quantum chemical study of two new inverse-sandwich complexes,  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$ . The global minimum structure of  $\text{La}_2\text{B}_{10}^-$  is found to be chiral ( $C_1$ ,  $^2A$ ) with an expanded inverse-sandwich structure, which can be viewed as adding a  $\text{B}_2$  unit off-plane parallelly to the  $\text{B}_8$  ring in a  $\text{La}_2\text{B}_8^-$  inverse-sandwich cluster. The global minimum of  $\text{La}_2\text{B}_{11}^-$  also has an expanded inverse-sandwich structure with a planar  $\text{B}_{11}$  framework, which can be viewed as adding three boron atoms to a  $\text{B}_8$  ring as the second coordination shell. Chemical bonding analyses show that the expanded  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$  inverse sandwiches have similar bonding as that in the monocyclic ring inverse sandwiches of  $\text{Ln}_2\text{B}_x^-$  ( $x = 7-9$ ), in which unique (d-p) $\delta$  bonding between the Ln...Ln atoms and the  $\pi$  orbitals on the boron rings was critical to stabilize the inverse-sandwich structures. The  $\text{La}_2\text{B}_{11}^-$  cluster is found to have an exceptionally stable electron configuration with a large HOMO-LUMO gap and very high electron binding energy. The current study confirms that the largest monocyclic boron ring to form inverse sandwiches in dilanthanide boride clusters ( $\text{Ln}_2\text{B}_x^-$ ) is nine. The  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$  clusters enrich the family of inverse-sandwich complexes and provide insight into the structural evolution of larger lanthanide boride clusters.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jpca.1c01149>.

Low-lying isomers of  $\text{La}_2\text{B}_{10}^-$  and  $\text{La}_2\text{B}_{11}^-$ , structural parameters and molecular orbitals of the global minima, comparison of simulated spectra, additional bonding analyses and theoretical VDEs, ADEs, and atomic charges, and Cartesian coordinates (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Authors

**Wan-Lu Li** – Department of Chemistry and Key Laboratory of Organic Optoelectronics & Molecular Engineering of Ministry of Education, Tsinghua University, Beijing 100084, China; Email: [wanyluli0716@gmail.com](mailto:wanyluli0716@gmail.com)

**Jun Li** – Department of Chemistry and Key Laboratory of Organic Optoelectronics & Molecular Engineering of Ministry of Education, Tsinghua University, Beijing 100084, China; Department of Chemistry, Southern University of Science and Technology, Shenzhen 518055, China; [orcid.org/0000-0002-8456-3980](https://orcid.org/0000-0002-8456-3980); Email: [junli@tsinghua.edu.cn](mailto:junli@tsinghua.edu.cn)

**Lai-Sheng Wang** – Department of Chemistry, Brown University, Providence, Rhode Island 02912, United States; [orcid.org/0000-0003-1816-5738](https://orcid.org/0000-0003-1816-5738); Email: [lai-sheng\\_wang@brown.edu](mailto:lai-sheng_wang@brown.edu)

### Authors

**Zhi-Yu Jiang** – Department of Chemistry and Key Laboratory of Organic Optoelectronics & Molecular Engineering of Ministry of Education, Tsinghua University, Beijing 100084, China

Teng-Teng Chen – Department of Chemistry, Brown University, Providence, Rhode Island 02912, United States; [orcid.org/0000-0001-9480-615X](https://orcid.org/0000-0001-9480-615X)

Wei-Jia Chen – Department of Chemistry, Brown University, Providence, Rhode Island 02912, United States

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.jpca.1c01149>

## Author Contributions

#Z.Y.J. and T.T.C. contributed equally to this work.

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

The experiment done at Brown University was supported by the National Science Foundation (CHE-1763380). The theoretical work done at Tsinghua University was supported by the National Natural Science Foundation of China (22033005). The calculations were done using supercomputers at the Southern University of Science and Technology (SUSTech) and Tsinghua National Laboratory for Information Science and Technology. The support of Guangdong Provincial Key Laboratory of Catalysis (No. 2020B121201002) is also acknowledged.

## REFERENCES

- (1) Nagamatsu, J.; Nakagawa, N.; Muranaka, T.; Zenitani, Y.; Akimitsu, J. Superconductivity at 39 K in magnesium diboride. *Nature* **2001**, *410*, 63–64.
- (2) Chung, H. Y.; Weinberger, M. B.; Levine, J. B.; Kavner, A.; Yang, J.-M.; Tolbert, S. H.; Kaner, R. B. Synthesis of ultra-incompressible superhard rhenium diboride at ambient pressure. *Science* **2007**, *316*, 436–439.
- (3) Kang, J. S.; Li, M.; Wu, H.; Nguyen, H.; Hu, Y. Experimental observation of high thermal conductivity in boron arsenide. *Science* **2018**, *361*, 575–578.
- (4) Li, S.; Zheng, Q.; Lv, Y.; Liu, X.; Wang, X.; Huang, P. Y.; Cahill, D. G.; Lv, B. High thermal conductivity in cubic boron arsenide crystals. *Science* **2018**, *361*, 579–581.
- (5) Tian, F.; Song, B.; Chen, X.; Ravichandran, N. K.; Lv, Y.; Chen, K.; Sullivan, S.; Kim, J.; Zhou, Y.; Liu, T.-H.; et al. Unusual high thermal conductivity in boron arsenide bulk crystals. *Science* **2018**, *361*, 582–585.
- (6) Tian, F.; Ren, Z. High thermal conductivity in boron arsenide: from prediction to reality. *Angew. Chem., Int. Ed.* **2019**, *58*, 5824–5831.
- (7) Carencio, S.; Portehault, D.; Boissiere, C.; Mezailles, N.; Sanchez, C. Nanoscaled metal borides and phosphides: recent developments and perspectives. *Chem. Rev.* **2013**, *113*, 7981–8065.
- (8) Sussardi, A.; Tanaka, T.; Khan, A. U.; Schlapbach, L.; Mori, T. Enhanced thermoelectric properties of samarium boride. *J. Materiomics* **2015**, *1*, 196–204.
- (9) Akopov, G.; Yeung, M. T.; Kaner, R. B. Rediscovering the crystal chemistry of borides. *Adv. Mater.* **2017**, *29*, 1604506.
- (10) Scheifers, J. P.; Zhang, Y.; Fokwa, B. P. T. Boron: enabling exciting metal-rich structures and magnetic properties. *Acc. Chem. Res.* **2017**, *50*, 2317–2325.
- (11) Alexandrova, A. N.; Boldyrev, A. I.; Zhai, H. J.; Wang, L. S. All-boron aromatic clusters as potential new inorganic ligands and building blocks in chemistry. *Coord. Chem. Rev.* **2006**, *250*, 2811–2866.
- (12) Wang, L. S. Photoelectron spectroscopy of size-selected boron clusters: from planar structures to borophenes and borospherenes. *Int. Rev. Phys. Chem.* **2016**, *35*, 69–142.
- (13) Li, W. L.; Chen, X.; Jian, T.; Chen, T. T.; Li, J.; Wang, L. S. From planar boron clusters to borophenes and metalloborophenes. *Nat. Rev. Chem.* **2017**, *1*, 0071.
- (14) Pan, S.; Barroso, J.; Jalife, S.; Heine, T.; Asmis, K. R.; Meirino, G. Fluxional boron clusters: from theory to reality. *Acc. Chem. Res.* **2019**, *52*, 2732–2744.
- (15) Jian, T.; Chen, X.; Li, S. D.; Boldyrev, A. I.; Li, J.; Wang, L. S. Probing the structures and bonding of size-selected boron and doped-boron clusters. *Chem. Soc. Rev.* **2019**, *48*, 3550–3591.
- (16) Kiran, B.; Bulusu, S.; Zhai, H. J.; Yoo, S.; Zeng, X. C.; Wang, L. S. Planar-to-tubular structural transition in boron clusters: B<sub>20</sub> as the embryo of single-walled boron nanotubes. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 961–964.
- (17) Oger, E.; Crawford, N. R.; Kelting, R.; Weis, P.; Kappes, M. M.; Ahlrichs, R. Boron cluster cations: transition from planar to cylindrical structures. *Angew. Chem., Int. Ed.* **2007**, *46*, 8503–8506.
- (18) Piazza, Z. A.; Hu, H. S.; Li, W. L.; Zhao, Y. F.; Li, J.; Wang, L. S. Planar hexagonal B<sub>36</sub> as a potential basis for extended single-atom layer boron sheets. *Nat. Commun.* **2014**, *5*, 3113.
- (19) Li, W. L.; Chen, Q.; Tian, W. J.; Bai, H.; Zhao, Y. F.; Hu, H. S.; Li, J.; Zhai, H. J.; Li, S. D.; Wang, L. S. The B<sub>35</sub> cluster with a double-hexagonal vacancy: a new and more flexible structural motif for borophene. *J. Am. Chem. Soc.* **2014**, *136*, 12257–12260.
- (20) Bai, H.; Chen, T. T.; Chen, Q.; Zhao, X. Y.; Zhang, Y. Y.; Chen, W. J.; Li, W. L.; Cheung, L. F.; Bai, B.; Cavanagh, J.; et al. Planar B<sub>41</sub><sup>-</sup> and B<sub>42</sub><sup>-</sup> clusters with double-hexagonal vacancies. *Nanoscale* **2019**, *11*, 23286–23295.
- (21) Zhai, H. J.; Zhao, Y. F.; Li, W. L.; Chen, Q.; Bai, H.; Hu, H. S.; Piazza, Z. A.; Tian, W. J.; Lu, H. G.; Wu, Y. B.; et al. Observation of an all-boron fullerene. *Nat. Chem.* **2014**, *6*, 727–731.
- (22) Chen, Q.; Li, W. L.; Zhao, Y. F.; Zhang, S. Y.; Hu, H. S.; Bai, H.; Li, H. R.; Tian, W. J.; Lu, H. G.; Zhai, H. J.; et al. Experimental and theoretical evidence of an axially chiral borospherene. *ACS Nano* **2015**, *9*, 754–760.
- (23) Romanescu, C.; Galeev, T. R.; Li, W. L.; Boldyrev, A. I.; Wang, L. S. Aromatic metal-centered monocyclic boron rings: Co@B<sub>8</sub><sup>-</sup> and Ru@B<sub>9</sub><sup>-</sup>. *Angew. Chem., Int. Ed.* **2011**, *50*, 9334–9337.
- (24) Galeev, T. R.; Romanescu, C.; Li, W. L.; Wang, L. S.; Boldyrev, A. I. Observation of the highest coordination number in planar species: decaordinated Ta@B<sub>10</sub><sup>-</sup> and Nb@B<sub>10</sub><sup>-</sup> anions. *Angew. Chem., Int. Ed.* **2012**, *51*, 2101–2105.
- (25) Romanescu, C.; Galeev, T. R.; Li, W. L.; Boldyrev, A. I.; Wang, L. S. Transition-metal-centered monocyclic boron wheel clusters (M@B<sub>n</sub>): a new class of aromatic borometallic compounds. *Acc. Chem. Res.* **2013**, *46*, 350–358.
- (26) Popov, I. A.; Jian, J.; Lopez, G. V.; Boldyrev, A. I.; Wang, L. S. Cobalt-centered boron molecular drums with the highest coordination number in the CoB<sub>16</sub><sup>-</sup> cluster. *Nat. Commun.* **2015**, *6*, 8654.
- (27) Jian, T.; Li, W. L.; Chen, X.; Chen, T. T.; Lopez, G. V.; Li, J.; Wang, L. S. Competition between drum and quasi-planar structures in RhB<sub>18</sub><sup>-</sup>: motifs for metallo-boron nanotubes and metallo-borophenes. *Chem. Sci.* **2016**, *7*, 7020–7027.
- (28) Jian, T.; Li, W. L.; Popov, I. A.; Lopez, G. V.; Chen, X.; Boldyrev, A. I.; Li, J.; Wang, L. S. Manganese-centered tubular boron cluster-MnB<sub>16</sub><sup>-</sup>: a new class of transition-metal molecules. *J. Chem. Phys.* **2016**, *144*, 154310.
- (29) Li, W. L.; Jian, T.; Chen, X.; Li, H. R.; Chen, T. T.; Luo, X. M.; Li, S. D.; Li, J.; Wang, L. S. Observation of a metal-centered B<sub>2</sub>-Ta@B<sub>18</sub><sup>-</sup> tubular molecular rotor and a perfect Ta@B<sub>20</sub><sup>-</sup> boron drum with the record coordination number of twenty. *Chem. Commun.* **2017**, *53*, 1587–1590.
- (30) Cheng, S.-B.; Berkdemir, C.; Castleman, A. Observation of d-p hybridized aromaticity in lanthanum-doped boron clusters. *Phys. Chem. Chem. Phys.* **2014**, *16*, 533–539.
- (31) Cheng, S.-B.; Berkdemir, C.; Castleman, A. W. Mimicking the magnetic properties of rare earth elements using superatoms. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112*, 4941–4945.
- (32) Chen, T. T.; Li, W. L.; Jian, T.; Chen, X.; Li, J.; Wang, L. S. PrB<sub>7</sub><sup>-</sup>: a praseodymium-doped boron cluster with a Pr<sup>II</sup> center



coordinated by a doubly aromatic planar  $\eta^7\text{-B}_7^{3-}$  Ligand. *Angew. Chem., Int. Ed.* **2017**, *56*, 6916–6920.

(33) Robinson, P. J.; Zhang, X.; McQueen, T.; Bowen, K. H.; Alexandrova, A. N.  $\text{SmB}_6^-$  cluster anion: covalency involving f orbitals. *J. Phys. Chem. A* **2017**, *121*, 1849–1854.

(34) Chen, X.; Chen, T. T.; Li, W. L.; Lu, J. B.; Zhao, L. J.; Jian, T.; Hu, H. S.; Wang, L. S.; Li, J. Lanthanides with unusually low oxidation states in the  $\text{PrB}_3^-$  and  $\text{PrB}_4^-$  boride clusters. *Inorg. Chem.* **2019**, *58*, 411–418.

(35) Mason, J. L.; Harb, H.; Huizenga, C. D.; Ewigleben, J. C.; Topolski, J. E.; Hratchian, H. P.; Jarrold, C. C. Electronic and molecular structures of the  $\text{CeB}_6$  monomer. *J. Phys. Chem. A* **2019**, *123*, 2040–2048.

(36) Chen, T. T.; Li, W. L.; Chen, W. J.; Li, J.; Wang, L. S.  $\text{La}_3\text{B}_{14}^-$ : an inverse triple-decker lanthanide boron cluster. *Chem. Commun.* **2019**, *55*, 7864–7867.

(37) Chen, T. T.; Li, W. L.; Chen, W. J.; Yu, X. H.; Dong, X. R.; Li, J.; Wang, L. S. Spherical trihedral metallo-borosphenes. *Nat. Commun.* **2020**, *11*, 2766.

(38) Li, W. L.; Chen, T. T.; Xing, D. H.; Chen, X.; Li, J.; Wang, L. S. Observation of highly stable and symmetric lanthanide octa-boron inverse sandwich complexes. *Proc. Natl. Acad. Sci. U. S. A.* **2018**, *115*, E6972–E6977.

(39) Chen, T. T.; Li, W. L.; Li, J.; Wang, L. S.  $[\text{La}(\eta^x\text{-B}_x)\text{La}]^-(x = 7-9)$ : a new class of inverse sandwich complexes. *Chem. Sci.* **2019**, *10*, 2534–2542.

(40) Li, W. L.; Chen, T. T.; Jiang, Z. Y.; Wang, L. S.; Li, J. Recent progresses in the investigation of rare-earth boron inverse sandwich clusters. *Chin. J. Struct. Chem.* **2020**, *39*, 1009–1018.

(41) Chen, X.; Zhao, Y. F.; Wang, L. S.; Li, J. Recent progresses of global minimum searches of nanoclusters with a constrained basin-hopping algorithm in the TGMin program. *Comput. Theor. Chem.* **2017**, *1107*, 57–65.

(42) Zhao, Y.; Chen, X.; Li, J. TGMin: A global-minimum structure search program based on a constrained basin-hopping algorithm. *Nano Res.* **2017**, *10*, 3407–3420.

(43) Chen, X.; Zhao, Y. F.; Zhang, Y. Y.; Li, J. TGMin: an efficient global minimum searching program for free and surface-supported clusters. *J. Comput. Chem.* **2019**, *40*, 1105–1112.

(44) Te Velde, G. T.; Bickelhaupt, F. M.; Baerends, E. J.; Fonseca Guerra, C.; van Gisbergen, S. J.; Snijders, J. G.; Ziegler, T. Chemistry with ADF. *J. Comput. Chem.* **2001**, *22*, 931–967.

(45) Guerra, C. F.; Snijders, J. G.; Te Velde, G. T.; Baerends, E. J. Towards an order-N DFT method. *Theor. Chem. Acc.* **1998**, *99*, 391–403.

(46) Baerends, E. J.; Ziegler, T.; Autschbach, J.; Bashford, D.; Bérces, A.; Bickelhaupt, F. M.; Yakovlev, A. L.; et al. *ADF 2017*; SCM, Inc.: Amsterdam, 2017.

(47) Perdew, J. P.; Burke, K.; Ernzerhof, M. Generalized gradient approximation made simple. *Phys. Rev. Lett.* **1996**, *77*, 3865–3868.

(48) Van Lenthe, E.; Baerends, E. J. Optimized Slater-type basis sets for the elements 1–118. *J. Comput. Chem.* **2003**, *24*, 1142–1156.

(49) Lenthe, E. v.; Baerends, E.-J.; Snijders, J. G. Relativistic regular two-component Hamiltonians. *J. Chem. Phys.* **1993**, *99*, 4597–4610.

(50) Adamo, C.; Barone, V. Toward reliable density functional methods without adjustable parameters: the PBE0 model. *J. Chem. Phys.* **1999**, *110*, 6158–6170.

(51) Dunning, T. H., Jr. Gaussian basis sets for use in correlated molecular calculations. I. The atoms boron through neon and hydrogen. *J. Chem. Phys.* **1989**, *90*, 1007–1023.

(52) Cao, X.; Dolg, M. Valence basis sets for relativistic energy-consistent small-core lanthanide pseudopotentials. *J. Chem. Phys.* **2001**, *115*, 7348–7355.

(53) Cao, X.; Dolg, M. Segmented contraction scheme for small-core lanthanide pseudopotential basis sets. *J. Mol. Struct.: THEOCHEM* **2002**, *581*, 139–147.

(54) Li, J.; Li, X.; Zhai, H. J.; Wang, L. S.  $\text{Au}_{20}$ : A tetrahedral cluster. *Science* **2003**, *299*, 864–867.

(55) Schipper, P. R. T.; Gritsenko, O. V.; van Gisbergen, S. J. A.; Baerends, E. J. Molecular calculations of excitation energies and (hyper) polarizabilities with a statistical average of orbital model exchange-correlation potentials. *J. Chem. Phys.* **2000**, *112*, 1344–1352.

(56) Zubarev, D. Y.; Boldyrev, A. I. Developing paradigms of chemical bonding: adaptive natural density partitioning. *Phys. Chem. Chem. Phys.* **2008**, *10*, 5207–5217.

(57) Mulliken, R. S. Electronic population analysis on LCAO-MO molecular wave functions. I. *J. Chem. Phys.* **1955**, *23*, 1833–1840.

(58) Hirshfeld, F. L. Bonded-atom fragments for describing molecular charge densities. *Theor. Chim. Acta* **1977**, *44*, 129–138.

(59) Wiberg, K. B.; Rablen, P. R. Comparison of atomic charges derived via different procedures. *J. Comput. Chem.* **1993**, *14*, 1504–1518.

(60) Bickelhaupt, F. M.; van Eikema Hommes, N. J. R.; Fonseca Guerra, C.; Baerends, E. J. The carbon-lithium electron pair bond in  $(\text{CH}_3\text{Li})_n$  ( $n = 1, 2, 4$ ). *Organometallics* **1996**, *15*, 2923–2931.

(61) Swart, M.; van Duijnen, P. T.; Snijders, J. G. A charge analysis derived from an atomic multipole expansion. *J. Comput. Chem.* **2001**, *22*, 79–88.

(62) Zhai, H. J.; Kiran, B.; Li, J.; Wang, L. S. Hydrocarbon analogs of boron clusters: Planarity, aromaticity, and antiaromaticity. *Nat. Mater.* **2003**, *2*, 827–833.

(63) Sergeeva, A. P.; Popov, I. A.; Piazza, Z. A.; Li, W. L.; Romanescu, C.; Wang, L. S.; Boldyrev, A. I. Understanding boron through size-selected clusters: structure, chemical bonding, and fluxionality. *Acc. Chem. Res.* **2014**, *47*, 1349–1358.

(64) Boldyrev, A. I.; Wang, L. S. Beyond organic chemistry: Aromaticity in atomic clusters. *Phys. Chem. Chem. Phys.* **2016**, *18*, 11589–11605.